

Incidence of Acute Transfusion Reaction and its Associated Risk Factors in Ateritiary Care Center in Ethiopia

Temesgen Assefa Ayele ^{1*}, Fissehatsion Tadesse ²

^{1*} Consultant Hematologist, Saint peter Hospital

² Consultant Hematologist, Assistant Professor of Hematology, Adissababa University

ARTICLE INFO

2024 Volume 1, Issue 1
<https://www.doi.org/jhrbd.2024.tgc.0264>

Article History:

Received: Jan 28, 2024
Accepted: Feb 03, 2024
Accepted: Feb 07, 2024

Citation: Temesgen A., Fissehatsion T. (2024). Incidence of Acute Transfusion Reaction and its Associated Risk Factors in Ateritiary Care Center in Ethiopia. *Journal of Hematological Research and Blood Disorders. The Geek Chronicles.* 1(1): 1-8

Copyright: © 2024 Temesgen A, this is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Keywords: blood transfusion, ATR, hemovigilance

ABSTRACT

Background: Transfusion reactions are adverse events associated with the transfusion of whole blood or one of its components. There is no institutional or national data in Ethiopia regarding the incidence of ATR & their risk factors. The aim of this study is to measure the incidence of acute transfusion reactions (ATR), frequency & time of occurrence of each ATR & evaluate their association with risk factors there by contributing to the institutional & national hemovigilance system.

Objective: The main objective of the study is to measure the incidence of ATR & determine their association with different clinical variables.

Methods: Prospective cross sectional study design is used to study in patient & outpatient adult transfusion recipients at the department of internal medicine, data was collected with a structured check list, analyzed with IBM SPSS version 24 software.

Result: A total of 210 study participants with the age of 13 & above were included in the study from September 1, 2022 – November 30 2022. 50.5 % of study participants are male & 49.5% are females, the most common age group are between the age of 18 -40, the commonest blood group was O positive. Acute myelogenous leukemia is the commonest underlying diagnosis of the study participants; PRBC is transfused for the majority of patients given for 74.8% of cases. ATR incidence is found to be 10%, ATR was reported in 19.6% of platelet transfused patients as compared to 7% of PRBC, and none of FFP transfused patients develop ATR. FNHTR & urticaria are the commonest ATR observed. Significant association is seen between ATR & female gender, autoimmune disease & ABO incompatible platelet transfusion.

Conclusion: The incidences of acute transfusion reactions is higher than reported from previous studies indicating the need to improve institutional transfusion services particularly in high-risk patients.

Introduction

A blood transfusion is an acute intervention, implemented to solve life and health-threatening conditions on a short-term basis by infusion of donated whole blood or blood components like RBCs, platelets, white cells & the different coagulation proteins [1]. Blood transfusion remains a common practice in the management of life-threatening clinical situations like trauma, surgical blood loss, severe anemia & different bleeding disorders to replace missing clotting factors and immune system elements [2,3]. Although Blood transfusion is a back bone for the management of a variety of medical & surgical patients, it is not without risks & complications.

Approximately 20% of all transfusions may lead to some type of adverse reactions. Complications associated with blood transfusion therapy may be classified based on time of onset as acute and late transfusion reactions or based on etiology as immunological and non-immunological.

Early onset transfusion reactions are usually acute reactions that occur during transfusion or anytime within 24 hours following transfusion of the blood or blood components, while late reactions occur from 24 hours to 2 weeks following the transfusion [4].

ATRs include acute hemolytic transfusion reaction (AHTR), allergic reactions, febrile non-hemolytic transfusion reaction (FNHTR), transfusion associated circulatory overload (TACO), and transfusion related acute lung injury (TRALI) and anaphylactic reactions [5].

The type and severity of transfusion reactions vary with the transfused blood product, the clinical condition of the recipient, past medical history and age of the recipient [6].

Despite the increasing public awareness on the risks and complications of blood transfusion that has resulted in a more stringent approach to donation, testing, and preparation of blood and its components, blood transfusion still is not without complications.

Thus, it is important for health professionals to monitor patients during and after transfusion &

it is essential to establish a system for monitoring, recording and reporting adverse reactions caused by blood transfusion in each hospital, thereby contributing to the national hemovigilance System [7].

Reports of blood component transfusion reactions are variable across the globe depending on the quality of the blood transfusion service at the center. The frequency of the complications is however inversely related with the care exercised in the preparation for and supervision of the transfusion [8]. In the developing countries blood transfusion services are fragmented, nonuniform, with different levels of care depending on the institution which will increase risk of ATRs. Reported incidences of ATR differ significantly while incidence of 0.2% and 0.34% are reported in Europe and South America, the incidence of acute immune-mediated transfusion reactions is reported to be 11.8% in North East Nigeria [9].

Learning from transfusion complications can drive the introduction of measures to enhance the quality, safety, efficacy and cost-effectiveness of blood and blood products as well as of the donation and transfusion processes [10]. The risk factors for common ATRs are reported only in few research papers & reviews performed before. In Ethiopia the incidence of ATRs & their associated risk factors is not known & there is no organized institutional hemovigilance system which record & report ATRs to the national blood bank. The aim of this study is to measure the incidence of patients who developed ATRs, frequency & time of occurrence of each ATR & evaluate their association with risk factors from patients transfused with blood components at medical wards, ICU & out patient hematology unit transfusion department of Tikuranbessa specialized hospital during the study period there by contributing to the safe blood transfusion practice & improving the blood transfusion surveillance system in the hospital.

Materials & Methods

3.1 Study design

A Cross-sectional prospective study was conducted from patients who were transfused with Blood components at medical wards, ICU & out patient transfusion department of Tikuranbessa specialized hospital during the study period.

3.2 Study period

The study was conducted from September 1, 2022 to November 30, 2022

3.3 Study area

The study was conducted at TASH which is located in Addis Ababa, the capital city of Ethiopia. It is the biggest referral hospital in the country with 700 inpatient beds and providing service to an estimated 500,000 patients annually. It also serves as a teaching hospital for undergraduate & post graduate medical & other health science students under the administration of Escanaba university college of health sciences.

3.4 Study population

All Patients who were transfused with different Blood components at medical wards, ICU & out patient transfusion department of TASH during the study period.

3.4.1 Inclusion criteria

- Age ≥ 13 years
- Patients who were transfused with blood products at Internal medicine department wards & medical ICU
- Patients who received blood component transfusion at outpatient hematology unit transfusion department.

3.4.2 Exclusion criteria

- Patients Age < 13 years
- Patients who received transfusion at hospital units other than Internal medicine Wards, medical ICU & outpatient hematology unit.

Result

4.1 Socio demographic characteristics of the study participants

From September 1 to November 30, 2022, a total of 210 patients who were transfused with different blood component during the study period were included in the study. 106 (50.5%) of patients are males & 104 (49.5%) are females. The study patients were categorized with the age group & the most common age group included in the study were between age group 18- 40 accounting for 92 (43.8%) of patients.




Figure 1: Age & sex distribution of study participants

4.2 Blood Group & diagnosis of study participants

The most common blood group identified among the study participants was O blood group which account for 40 % of the study population, others include blood group B (27.2%), A (26.2%) & AB (6.7%) of patients. The RH status of the study also studied & showed 198 (94.3%) were RH positive & 12 (5.7%) were RH negative. Considering both the

blood group & Rh status of the patients O+ve is the commonest blood group seen in 79 (37.6%) of patients.

The three most common medical problems identified among the study participants were Acute myelogenous leukemia, Aplastic anemia & Chronic myelogenous leukemia which are observed in 62 (29.5%), 32 (15.2%) & 24 (11.4%) patients respectively.




Figure 2: Blood group of study participants

4.3 Type of blood components transfused

Blood components transfused for the study population were Packed RBC, Platelets & FFP which were given for 157 (74.8%), 51(24.3%), 2 (1 %) patients respectively.

4.4 Proportion of Acute Transfusion reaction

From the total of 210 transfused patients with different blood component products 21 (10%) patients develop acute transfusion reaction during the study period; 11 of which are seen among patients transfused with PRBC, 10 patients develop ATR after platelet transfusion. Based on patients' subjective report of ATR 32 (15.2%) of patients developed at least one episode of ATR previously.

4.5. Type of ATR

FNHTR & urticaria are the only acute complications of transfusion identified during the study period which were seen in 12 (5.7%) & 9 (4.3 %) patients respectively.

When ATR were analyzed according to blood components, ATR was seen in 11 (7%) patients transfused with packed RBC & 10 (19.6%) patients transfused with platelets, no patients transfused with FFP developed ATR.

4.6 Time of occurrence of ATR

The time of occurrence of ATR was variable among study subjects. The majority of patients (52.4%) with ATR develop within a period of one to six hours.

Table 1 - Time of occurrence of ATR

Time of occurrence of ATR	Frequency	Percent
< 30 minutes	3	14.3 %
30 minutes - 1 hour	7	33.3%
1 hour – 6 hours	11	52.4%

4.7 Association of ATR with different clinical variables

4.7.1 ATR & Gender

From patients who developed ATR 15 (71.6%) were females & 6(28.6%) were males. There is significant association between the patients' gender & the development of ATR with two-sided significance **P value of 0.04** by Pearson chi-square & Fischer's exact test.

4.7.2 ATR & Age

ATR developed most commonly in age group 18 - 40 years accounting for 47.6 % of ATR cases but there was no significant association between age group differences in the development of ATR with Pearson chi square test **P value of 0.21**.

4.7.2 ATR & Blood group

There was no significant association between the blood group of the study subjects & the

development of ATR with Pearson chi square test **p value of 0.09**.

4.7.3 ATR & blood component transfused

Among 157 patients transfused with Packed RBC only 7% of patients developed ATR but 19.6 % of patients among 51 platelet transfused patients developed ATR. Large number of patients who are transfused with platelets developed ATR than those transfused with PRBC & FFP & there is a significant association between type of blood component transfused & the development of ATR with a Pearson chi square **P value of 0.03**.

4.7.4 ATR & Previous history of transfusion

Out of the 32 patients with previous history of transfusion 8 patients developed ATR & there is no significant association observed between current ATR & previous history of transfusion with **2-sided significance level of 0.06 (P >0.05)** by Pearson chi square test.

Table 2 – Distribution of ATR by previous transfusion

Acute transfusion reaction	Previous transfusion		Total
	Yes	No	
Yes	8	13	21
No	24	165	189

P = 0.06 (Fisher exact) Chi-square = 9.438 df = 1

5.7.5 ATR & autoimmune disease

11 patients (5.7 %) of the total study population were found to have autoimmune diseases. From those who developed ATR 4 (19%) patients has autoimmune disease documented & the association between ATR & the presence of autoimmune disease is observed with Pearson chi square **P value of 0.011**.

4.7.6 ATR & ABO incompatibility

ABO incompatible transfusion was seen in 6 (2.9%) of the study patients & all were observed during platelet transfusion. ABO mismatch transfusion is seen in 60 % of platelet transfused patients who developed ATR.

Significant association observed between the presence of ABO incompatibility & ATR with Chi square lambda correlational analysis **P value of 0.013**.

Discussion

A total of 210 blood recipients participated in this study, out of which 106 (50.5 %) of patients are males & 104 (49.5%) are females with M; F ratio of 1.1: 1 & the commonest age group reported to receive blood component transfusion during the study period are the age group between 18- 40.

The commonest diagnosis reported among transfusion recipients is acute myelogenous leukemia which accounts for 29.5% of the study subjects which is different from the study reported in Nigeria by Baffa A. et al which reported the commonest diagnosis among the blood recipients was HIV/AIDS accounting for 16.1% cases while malignancies in general accounted for 15.6% cases; [8] this may be due to the difference in the baseline characteristics of the study population ,our study is performed in adult medical patients in a tertiary care hospital where majority of transfused patients have the diagnosis of hematologic malignancies. ATR is seen in 10 % of our study population which is higher than the studies done at U.S which reported ATR incidence rate of 0.2% in the study reported by Rohi et al, it is also higher than the studies reported in Namibia, Nigeria which reported incidence rate of 0.2%,3.6% & also higher than the study done in other area of Ethiopia which reported ATR rate of 5.2% (20 patients out of 364 blood recipients) [8-14].

In our study the ATRs identified are only FNHTR & urticaria (acute allergic reactions) which are also the most common ATRs reported from different centers; FNHTR is seen in 57.1 % & urticaria in 42.9% among patients who developed ATR which is similar to the study by Pahuja et al which reported the frequency of FNHTR & urticaria as 54.7% & 41.4% respectively, [24] but it is different from the study by Yemataw et al which reported allergic reactions in 65% while FNHTR in 30% of ATRs [15].

Frequency of ATR is variable depending on the type of blood components transfused, in our study proportion of ATR is higher among platelet transfused patients which is seen in 19.6

% of 51 platelet transfused patients but only 7% of RBC transfused patients developed ATR. This finding is higher than the study done in Japan by Yuki Hatayama et al which reported ATR rate in 5.7% of platelet concentrate ,1.6% of red blood cell component and 2.2% FFP.

In study done in India by Niga et al ATR were seen mostly with whole blood (43.5%) and PRBCs (48.5%), ATR to platelets was seen in 6/101 (5.9%) cases [18, 23].

We evaluated the time from initiation of transfusion to the occurrence of signs and symptoms of ATR, the majority of ATR (52.4%) developed during the period of one to six hours after the initiation of transfusion & there was no significant difference between the time of occurrence of ATR, blood component transfused & type of ATR which is similar to the result in the study done in Japan by Yuki Hatayama et al. This indicates that patients need close follow-up during the first six hours after transfusion initiation [18].

In this study, a statistically significant relationship was obtained between ATR and the gender of the transfused patients, type of blood component transfused & the presence of autoimmune disease in which female, ABO mismatched platelet transfused & those with autoimmune diseases are associated with the development of ATRs ; this findings are different from studies performed in Nigeria which reported a significant relation between ATR with previous transfusion & the age of stored blood components [8];it is also different from the study by Yemata et al which reported ATR was associated with transfusion history ,abortion history, storage time and transfusion of 3 or more of units of blood/blood components [15].

ABO & RH incompatible platelet transfusion service is given by many centers due to inventory constraints when single donor apheresis platelets are not available; clinically significant transfusion reactions, though uncommon, are seen with ABO mismatched platelet transfusions [25].

In our study all (6 out of 45 platelet transfusions) of ABO mismatched platelet

transfusions are associated with ATR particularly acute allergic reactions (urticaria). In our study there is no significant association seen between ATR & the age, blood group, previous transfusion & previous ATR history of the study participants.

Conclusion

The incidence of ATR is 10 % of the studied population which is higher than the reports from other studies. FNHTR is the commonest ATR seen in 5.7 % & urticaria was seen in 4.3 % of the study subjects. ATR is seen commonly in platelet transfused patients as compared to PRBC or FFP recipients. Significant association is seen between ATR & female gender, the presence of auto immune disease & ABO incompatible platelet transfusion unlike other studies in which significant association of ATR was seen with previous transfusion history, previous abortion history & baseline leukopenia & storage time of the blood product.

The commonest time of occurrence of ATR is the first six hours after initiation of the transfusion.

It is recommended from this study that Blood transfusion is not without risks, transfused patients need due attention & close follow-up particularly during the first six hours of transfusion. Female patients especially those with autoimmune disease need especial attention due to the strong association with the development of ATR.

Platelet transfused patients need to be followed carefully for the development of ATR & ABO mismatched platelet transfusion need to be

avoided as much as possible by advocating single donor platelet transfusion practice.

Strengths and Limitations of the Study

This study is one of the few studies performed on transfusion medicine & the 2nd of its kind reporting the incidence of ATR & associated factors in the country so far. It is a prospective cross-sectional study which tries to assess the incidence of ATR, associated factors & their occurrence time in a setup where there is no strong transfusion reaction reporting system which will help to contribute data to the development of institutional & national hemovigilance system.

Small sample size, short study period & the limited group of patients studied are the limitations of the study which will make the generalization of the results to the wider transfusion recipient population difficult; therefore, it is recommended to study wider population over a longer period of observation in the future so that the incidence of ATRs including those that are not identified in our study will be determined.

Conflict of Interest

No conflict of Interest

Authors Contribution

Temesgen Assefa Ayele is a corresponding author involved in design of the study, data collection, analysis & write up of the study.

Fissehatsion Tadesse is coauthor involved in the design & analysis of the study.

References

1. WHO Clinical Transfusion Practice Guideline
2. Jeffrey JM. Transfusion Medicine. 3rd ed,2012.
3. Rao G HR, Eastlund T, Jagannathan L. Handbook of Blood Banking and Transfusion Medicine,2006
4. Walker RH. Special report: Transfusion risk. Am J Clin Pathol 1987; 88:374
5. Arewa OP. Evaluation of transfusion pyrexia: a review of differential diagnosis and management. ISRN Hematol. 2012; 2012:1–7.
6. Organization WH. Clinical transfusion practice: guidelines for medical interns. Clin Transfus Pract Guidel Med Interns. 2012:1–42
7. Yuki Hatayama et al, Analysis of Acute Transfusion Reactions and Their Occurrence Times, Yonago Acta Medica 2018
8. Baffa A. Gwaram, Musa M. Borodo, Abdulhamid I. Dutse, Aisha Kuliya-Gwarzo, Pattern of acute blood transfusion reaction in Kano, North western Nigeria
9. Ahmed SG, Ibrahim UA, Gamas MG et al, Incidence and clinical pattern of immune mediated blood transfusion reactions in Maiduguri, Nigeria
10. A guide to establishing a national haemovigilance system, World Health Organization 2016
11. Robert Weinstein, MD, A pocket guide for RBC Transfusion, University of Massachusetts Medical School, November 2016
12. Serious Hazards of Transfusion 2020 report in U.K
13. Rohit, Hindu etal, Incidence and Outcomes of Acute Transfusion Reactions in Hospitalized Patient in the United States, Blood 2020
14. Benjamin, Britta etal, Estimation of the prevalence and rate of acute transfusion reactions occurring in Windhoek, Namibia
15. Yemataw, Berhanu etal, Proportion of Acute Transfusion Reaction and Associated Factors Among Adult Transfused Patients at Felege Hiwot Compressive Referral Hospital, Bahir Dar, Northwest Ethiopia: A Cross-Sectional Study, Journal of Blood medicine
16. Yuki Hatayama et al, Analysis of Acute Transfusion Reactions and Their Occurrence Times, Yonago Acta Medica 2018
17. Chao-Yuan Yao et al, Associated Factors with Acute Transfusion Reaction from Hospital Online Reporting Events: A Retrospective Cohort Study, Journal of patient safety 2020
18. Yutki, Satoko et al, Analysis of Acute Transfusion Reactions and Their Occurrence Times, Yonago Acta Medica 2018
19. Chang, Lee etal, Transfusion-associated adverse reactions (TAARs) and cytokine accumulations in the stored blood components: the impact of prestorage versus poststorage leukoreduction.
20. Arthur, Steven et al, Uptodate 2018, Immunologic transfusion reactions
21. Alam, Lin et al, The prevention of transfusion-associated circulatory overload, Transfusion Medicine Review 2013
22. Vlaar, Binnekade et al, Risk factors and outcome of transfusion-related acute lung injury in the critically ill: a nested case-control study.
23. Negi, et al.: Adverse transfusion reactions, Advanced Biomedical Research
24. Pahuja S, Puri V, Mahajan G, Gupta P, Jain M. (2017). Reporting adverse transfusion reactions: A retrospective study from tertiary care hospital from New Delhi, India. Asian J Transfus Sci 11:6-12
25. Josephson et al. (2004). Significant numbers of apheresis-derived group O platelet units have "high-titer" anti-A/A, B: implications for transfusion policy. Transfusion 44:805.